

Biological and Biochemical Effects of *Bacillus thuringiensis*, *Serratia marcescens* and Teflubenzuron and their Sequential Combined Effects on Cotton Leafworm, *Spodoptera littoralis* (Boisd.).

El-Aasar A.M.¹; El-Sheikh, T.A.A.²; Rafea, Heba S.² and Ali S. H.¹.

1- Faculty of Agriculture , Ain –Shams- University, Egypt.

2- Plant Prot. Res. Inst., Agric Res. Centre, Dokki, Giza, Egypt.

ABSTRACT

Two biopesticides *Serratia marcescens* [Eubacteriales: *Enterobacteria*] (used at MC₅₀, concentration caused 50% malformation)and *Bacillus thuringiensis* Var. *kurstaki* (used at LC₅₀) and insect growth regulator Teflubenzuron (used at LC₅₀ value) were used for treatment of 2nd instar larvae of cotton leafworm, *Spodoptera littoralis* (Boisd.). Sequential combined Effect was carried out by treating 2nd instar larvae with LC₅₀ value of *B. thuringiensis* or Teflubenzuron then the larvae allowed to pupate on sawdust treated with *S. marcescens* at MC₅₀. The effects of these three agents were assessed by toxicity, fecundity, fertility, phosphatases activity and total carbohydrates. The obtained results revealed that Teflubenzuron is a potent toxin (LC₅₀ = 0.113 ppm) compared to *B. thuringiensis* (LC₅₀ = 165.64 ppm). On the other hand, the mode of action of *S. marcescens* is through malformation (MC₅₀ = 3.09x10⁸ cfu). All treatments caused a significant reduction in fecundity and fertility. The sequential combined effect showed more reducing effect on the fecundity and fertility than the individual treatments. Moreover, the activity of phosphatases and total carbohydrates were significantly fluctuated during the different periods of pupal stage. The sequential combined effect of Teflubenzuron with *S. marcescens* caused more significant effect on alkaline phosphatase activity than the individual treatment with *S. marcescens* but it was somewhat similar to Teflubenzuron. Also the sequential combined effect treatment of *B. thuringiensis* with *S. marcescens* had more effect than the individual treatment either with *S. marcescens* or *B. thuringiensis*. Moreover, all treatments caused significant decreases in total carbohydrates during the pupal stage and the sequential combined effect treatments had more decreasing effect than the individual treatments. According to the obtained result, *S. marcescens* could be considered as a biopesticide, and become more effective when used in sequential treatment with either Teflubenzuron or *B. thuringiensis*.

Key words: *Serratia marcescens*, *Bacillus thuringiensis* Var. *kurstaki*. Teflubenzuron, *Spodoptera littoralis*, biological, biochemical aspects.

INTRODUCTION

The cotton leafworm, *Spodoptera littoralis* (Boisd.) is a highly destructive insect pest. The extensive use of insecticides to control *S. littoralis* larvae has led to several problems as development of resistance and residual effects (Frank *et al.*, 1990). Thus, attention was directed to search for

alternative control agents with new modes of action. Among these agents are insect growth regulators (IGR's) and microbial control agents. IGR's have been classified as "bio-rational" to distinguish them from conventional insecticides. These compounds are selective and specific to the target pest (Metcalf *et al.*, 1975). Moreover, they

elicit their primary action on insect metabolism and ultimately affecting development and growth of the target insect. They induce morphological abnormalities as well as death of treated insects. These characteristics made them most promising as new control agents for controlling *S. littoralis* larvae. On the other hand, the bacterium *B. thuringiensis*, proved to be a highly successful weapon for fighting some agricultural pests and it offer many advantages over chemical insecticides. *B. thuringiensis* is known to be one of the most pathogenic species of bacteria, which induce larval mortality after a course of infection stages. The interest of using such agent as a microbial bio-insecticide was increased since 1970 (Dulmage and Co-operators, 1981). The bacteria of the genus *Serratia* are often associated with insects and have the behavior of a facultative pathogen. (Trevor *et al.*, 2004). In general, *Serratia marcescens* is not pathogenic to insects when present in the digestive tract in small numbers, but once it enters the hemocoel it multiplies rapidly and causes death in one to three days (Sikorowski, 1985). Furthermore, chitinase producing bacteria *S. marcescens* caused significant physiological and morphological effects on pupal and adult stages where it caused a significant increase in the proportion of pupal mortality, adult malformation and sterility with treated moths and also affected some enzymes activity (Tolba, 2006). The objective of this work is directed to study the assessment of *S. marcescens* as an biopesticide against *S. littoralis* with comparison to *B. thuringiensis* Var. *kurstaki* and the insect growth regulator Teflubenzuron as a chitin synthesis inhibitor and evaluation of sequential combined effect of *S. marcescens* with either *B. thuringiensis* or Teflubenzuron for controlling *S. littoralis*.

MATERIALS AND METHODS

Rearing technique

The stock culture of the cotton leafworm, *Spodoptera littoralis* (Boisd.) was obtained from a laboratory strain maintained for several generations without any insecticidal or microbial pressure, in the Cotton leafworm Research Dept, Plant Protection Research Institute, Agricultural Research Centre, Dokki, Giza. The insect was reared on castor-oil leaves, *Ricinus communis*; under laboratory conditions at 25 ± 2 °C and 60 ± 5 % R.H. 2nd and late 6th instars larvae were used in the current work.

Control agents

Biopesticide

Protecto: It is a wettable powder formulation, based on *Bacillus thuringiensis* Var. *kurstaki*. It contains lepidopteran toxin 9.4 % produced by the Plant Protection Research Institute, Agricultural Research Centre, Dokki, Giza, Egypt.

***Serratia marcescens*:** Chitinase-producing bacterial strain belongs to *Enterobacteriaceae* isolated from Egyptian Soils. The isolated bacterial strain was formulated as a biocontrol agent for controlling parasitic nematodes. It was produced by Soils, Water and Environ. Res. Inst. Agriculture research center, and distributed on a commercial scale (trade name, Nemaless)

IGR

Common name: Teflubenzuron.

Trade name: No moult 15 % S.C.

This IGR has been obtained from Sumitomo Chemical Co., Ltd., Japan.

Bioassay trials

Preliminary tests for the individual treatments were carried out using series of concentrations (in water) for each of the bio-agent, *Serratia marcescens* (10^5 , 10^6 , 10^7 , 10^8 , 10^9) colony forming unit/ml (cfu/ml), *B. thuringiensis* (44.187, 88.375, 176.75, 352.5, 705, 1410 ppm) and the chitin synthesis inhibitor

Teflubenzuron (0.02, 0.04, 0.08, 0.16, 0.32, 0.64 ppm). Sawdust was treated with each concentration of *Serratia marcescens* in glass jars and offered to late 6th instar larvae to pupate on it. The offered treated sawdust was in a wettable form, while, in case of Teflubenzuron and *B. thuringiensis*, the use of leaf-dipping technique was carried out according to Abo El-Ghar *et al.*, 1994. Castor bean leaves, *R. communis*, were dipped in each concentration then left to dry at room temperature and these were offered to the newly moulted 2nd instar larvae. Larvae were allowed to feed for 24 hrs. then, they were provided with fresh, clean and untreated castor bean leaves until pupation. Larvae that fed on untreated castor bean leaves were used as control for Teflubenzuron and *B. thuringiensis* treatments whereas larvae kept in untreated sawdust were considered as control for *Serratia marcescens*. In all treatments, three replicates were carried out for each concentration; each replicate consisted of 20 larvae. The larval mortality and adult malformation percentages were determined. The data were then subjected to probit analysis (Finney, 1971) to obtain the LC₅₀ values of both Teflubenzuron and *B. thuringiensis* as well as the concentration which causes 50% adult malformation (MC₅₀) for *S. marcescens*. The sequential combined effect treatments of *S. marcescens* either with *B. thuringiensis* (*Bt/Serr*) or

Teflubenzuron (*Teflu /Serr*) were carried out by treatment of the 2nd instar larvae with the obtained LC₅₀ either of *B. thuringiensis* or Teflubenzuron then, at the end of larval stage, the late 6th instar larvae were allowed to pupate on sawdust treated with *S. marcescens* at the MC₅₀. The Toxicity index of the tested compounds was determined according to Sun (1950) as follows:

$$\text{Toxicity index} = \frac{\text{LC}_{50} \text{ of the most toxic compound}}{\text{LC}_{50} \text{ of other compounds}} \times 100$$

Biological studies:

In all treatments, either the individual or the sequential treatments, the larvae were examined daily. The number of eggs deposited by mated female moth and the percentage of egg hatchability were recorded. Mating processes occurred according to the following combinations: Treated female x treated male, Treated female x untreated male, untreated female x Treated male and untreated female x untreated male (control). Each mating combination consisted of newly emerged of ratio one male to one female. Three replicates were carried out for each mating combination. Fecundity and fertility of *Spodoptera littoralis* moths in each treatment were determined. Fecundity was measured as the total number of eggs laid/female. The deterrent index based on the number of eggs in treatment and control assays was calculated according to Lundgren (1975) as follows:

$$\text{Deterrent index} = \frac{A - B}{A + B} \times 100$$

Where; A = number of eggs/female in control and B = number of

eggs/female in treatment. The egg hatchability % (fertility) was determined.

$$\text{Deterrent index} = \frac{A - B}{A + B} \times 100$$

**Biochemical Analysis
Preparation of samples for
biochemical analysis:**

Pupal homogenate samples were collected after 2, 4, 6, 8, 10, 12 and 14 days of pre-pupation and were homogenized in distilled water using a

Teflon homogenizer surrounded with a jacket of crushed ice for 3 minutes. Homogenates were centrifuged at 6000 r.p.m. for 10 minutes at 5°C. The supernatant was divided into small aliquots (0.5 ml) and stored at -20 °C until analysis. Three replicates were used for each biochemical determination.

Determination of phosphatases activity

Acid and alkaline phosphatase activities were measured according to the method of Laufer and Schin (1971).

Determination of total carbohydrates

Total carbohydrates were determined as described by Singh and Sinha (1977).

Statistical analysis

Data were analyzed by ANOVA using SAS package.

RESULTS AND DISCUSSION

Toxicological effects:

Data in Table (1) showed the susceptibility of the 2nd instar larvae of *S. littoralis* towards *B. thuringiensis* and Teflubenzuron compounds. The LC₅₀ of Teflubenzuron is 0.113ppm, whereas, it is 165.64ppm in case of *B. thuringiensis*. Based on LC₅₀ values, it is obvious that both compounds caused considerable toxic effects against the 2nd instar larvae

of *Spodoptera littoralis* particularly in case of Teflubenzuron which had drastic toxic effects comparing to *B. thuringiensis* toxicity. On the other hand, *S. marcescens* had no toxic effects against larvae. However it has the ability to cause a high malformation percentage to *Spodoptera littoralis*. The concentration which causes 50 % adult malformation is 3.09x10⁸ cfu.

Table 1: Toxicity of *B.thuringiensis* , teflubenzuron and MC₅₀ of *Serratia marcescens* to *Spodoptera littoralis*.

Agent	LC ₅₀ (ppm)	95% Fiducial Limits		Slope ± S.E.	X ² _(df)	Toxicity Index
<i>B.thuringiensis</i>	165.64	128.21	208.31	1.59 ± 0.12	0.523 (5)	0.068
Teflubenzuron	0.113	0.075	0.171	1.59 ± 0.13	9.94 (5)	100
<i>Serratia marcescens</i> concentration which causes 50% adult malformation to <i>spodoptera littoralis</i> (Boisd).						
Agent	*MC ₅₀ ** (cfu)	95% Fiducial Limits		Slope ± S.E.	X ²	Toxicity Index
<i>Serratia marcescens</i>	3.09x10 ⁸	7.7x10 ⁷	3.81x10 ⁹	0.29 ± 0.055	0.966 (4)	-

* MC₅₀ :Concentration caused 50% adult malformation.

** (cfu) : colony forming unite .

The toxicity of *B. thuringiensis* to *S. littoralis* larvae is due to the production of crystalline δ-endotoxin (cry) protein during the sporulation stage of *B. thuringiensis* life cycle. On digestion by susceptible larvae, the active toxin generated from the protoxin binds to the receptors on gut epithelium. This leads to paralysis of gut and mouth parts causing death of larvae (Aronson *et al.* 1986). On the other hand, Teflubenzuron, a potent chitin synthesis inhibitor classified as insect growth regulator,

inhibits the synthesis of chitin in larvae which have ingested it; causing the integument to become fragile, and leading to mortality during the moulting (Ascher and Nemny, 1984). *S. marcescens* as chitinase producing bacteria is one of the most effective bacteria for degradation of chitin. *S. marcescens* chitinases may find applications as biocontrol agents against insects Brurberg *et al.* (2000). This might explain the high malformation percentage

in *S.littoralis* treated with *S. marcescens* obtained in the present work.

Biological effects.

Fecundity:

Data in Table (2) indicated the changes in the fecundity (number of eggs laid per female) and oviposition deterrent index (OVDI) of *spodoptera littoralis*. The obtained results showed that, in all treatments, the highest reduction in the

fecundity, together with the highest oviposition deterrent index were obtained for treated female mated with treated males, followed by treated females mated with normal males. On the other hand, the highest fecundity and consequently the lowest OVDI were obtained when the males were only treated.

Table 2: Effect of treatment of 2nd instar larvae of *spodoptera littoralis* with LC₅₀ levels of *B.thuringiensis*, teflubenzuron and concentration which causes 50 % adult malformation of *S. marcescence* and their combined effect on the No. of eggs/female (fecundity) and oviposition deterrent index (OVDI) (*mean values ± SE).

Compound	Type of mating		Eggs / ♀ Mean ± SE		**OVDI	Compound	Type of mating		Eggs / ♀ Mean ± SE		OVDI
	♂	♀					♂	♀			
<i>Serr.</i>	Treated	Untreated	1062.67	c ± 36.08	22.98	<i>Bt / Serr.</i>	Treated	Untreated	992.67	c ± 31.05	26.18
	Untreated	Treated	910.33	d ± 28.86	30.16		Untreated	Treated	867	d ± 26.49	32.36
	Treated	Treated	695.33	e ± 19.12	41.86		Treated	Treated	685.33	e ± 24.07	42.46
<i>Bt</i>	Treated	Untreated	1160.33	b ± 34.71	18.77	<i>Teflu/Serr.</i>	Treated	Untreated	311.33	g ± 10.82	68.99
	Untreated	Treated	1040.33	c ± 28.85	23.98		Untreated	Treated	164.33	hi ± 5.24	82.34
	Treated	Treated	884.33	d ± 20.72	31.47		Treated	Treated	99	i ± 4.73	88.97
Teflubenzuron	Treated	Untreated	545.67	f ± 16.85	51.33	Control	Untreated	Untreated	1696.7	a ± 44.72	-
	Untreated	Treated	333.33	g ± 9.34	67.16						
	Treated	Treated	233.33	h ± 7.06	75.82						
LSD	71.205										

*Means with the same letter(s) are not significantly different.

**OVDI: Oviposition deterrent index

This may indicate that the females were more sensitive to the tested compounds than the males. Moreover, in all mating combinations of all treatments, the fecundity of the resulting treated females was significantly decreased as compared to control. In The individual treatments, Teflubenzuron had higher reducing effect on the fecundity particularly for treated female mated with treated males (OVDI =75.82) than the individual treatment either with *S. marcescens* (OVDI = 41.86) or *B. thuringiensis* which had the less reducing effect (OVDI = 31.47). Therefore, *S. marcescence* as a biopesticide is more effective on the fecundity than *B.*

thuringiensis but it less effective than Teflubenzuron . Moreover, the sequential combined effect treatments had higher reducing effect than the individual treatments where, Teflu /*Serr* treatment caused higher significant decreases in the fecundity of all combinations than the individual treatments with both *S. marcescens* and Teflubenzuron particularly for treated female mated with treated males. Also *Bt/Serr* treatment had reducing effect which was higher than the individual *B. thuringiensis* but it was similar to the individual treatment with *S. marcescens*. As compared to their control, it is clear that *S. marcescens* clearly enhances the reducing effect

either of *B. thuringiensis* or Teflubenzuron on *S. littoralis* fecundity.

The reduction in the fecundity of *S. littoralis* treated with *S. marcescens* was similar to that obtained by Tolba (2006) who found that *S. marcescens* significantly reduced the fecundity of *A. ipsilon*. In addition Abd El-Hameed (1995) found a significant reduction in fecundity of the *Pectinophora gossypiella* after treatment with *B. thuringiensis*. Moreover, Abdel-Aal and Abdel-Khalek (2006) found that Teflubenzuron significantly decreased the fecundity percentage of *S. littoralis* especially when treated females mated with treated males.

Fertility%

Data in Table (3) showed changes in the fertility % (eggs hatchability %) of *S. littoralis*. The obtained results revealed that all treatments in all mating combinations cause a significant decrease in fertility of *S. littoralis* as compared to control. The fertility followed the same pattern of that of fecundity (the lowest fertility percent was obtained in mating combinations contained treated females), emphasizing that the females were more sensitive to the tested compounds than the males.

The highest reduction in fertility was obvious in case of the sequential combined effect treatment Teflu /*Serr*. The reduction effects in the fertility were follow the sequence Teflu /*Serr* > Teflubenzuron > *S. marcescens* > *Bt/Serr* > *B. thuringiensis*, where, the fertility reduced to 15.15, 22.43, 35.23, 45.23 and 57.33 % when treated female mated with treated male, respectively. While, the reduction effects in the fertility were follow the sequence Teflu /*Serr* > Teflubenzuron > *Bt/Serr* > *S. marcescens* > *B. thuringiensis*, where, the fertility reduced to 39.55, 47.70, 55.52, 59.58 and 69.56 % when treated female mated with untreated male, respectively, as compared with control (the fertility being 98.25%). These results reflect the higher reducing effect of *S. marcescens* as a biopesticide on the fertility than *B. thuringiensis* whereas Teflubenzuron had higher reducing effect on the fertility than *B. thuringiensis* and *S. marcescens*. Also, it is clearly appeared the synergistic effects of *S. marcescens* in the sequential treatment either with Teflubenzuron or *B. thuringiensis* on the fertility of *S. littoralis*.

Table 3: Effect of treatment of 2nd instar larvae of *spodoptera littoralis* with LC₅₀ levels of *B.thuringiensis*, teflubenzuron and concentration which causes 50% adult malformation of *S. marcescens* and their combined effect on the egg hatchability %.

Type of mating		<i>S. marcescens</i>		<i>B. thuringiensis</i>		Teflubenzuron		<i>BT/SERR</i>		Teflu/ <i>SERR</i>	
?	?	Number of eggs *Mean ± SE	hatchability %	Number of eggs *Mean ± SE	hatchability %	Number of eggs *Mean ± SE	hatchability %	Number of eggs *Mean ± SE	hatchability %	Number of eggs *Mean ± SE	hatchability %
Treated	Untreated	686.00 ± 21.10 ^c	64.55	870.33 ± 21.34 ^b	75.01	290.67 ± 10.15 ^{gh}	53.27	597.67 ± 18.70 ^d	60.21	147.00 ± 5.70 ⁱ	47.22
Untreated	Treated	542.33 ± 15.08 ^e	59.58	723.67 ± 16.21 ^c	69.56	159.00 ± 7.78 ⁱ	47.70	481.33 ± 19.62 ^f	55.52	65.00 ± 4.73 ^j	39.55
Treated	Treated	245.00 ± 6.44 ^h	35.23	507.00 ± 14.31 ^{ef}	57.33	52.33 ± 4.67 ^{jk}	22.43	310 ± 13.24 ^g	45.23	15.00 ± 3.22 ^k	15.15
control (Untreated X Untreated)		1667 ± 44.49 ^a	98.25	-	-	-	-	-	-	-	-
LSD		49.713									

*Means with the same letter(s) are not significantly different.

Desneux *et al.* (2007) reported that reduction in the fecundity associated with pesticides may be due to both physiological and behavioral effects. The authors suggested also that insect growth regulators such as Teflubenzuron may induce mere long-term effects on fecundity than, for example, neurotoxins. Whereas, Haseeb and Amano (2002) recorded reduction in parasitism, when Teflubenzuron investigated by *Cotesia plutella* adults and suggested that this insect growth regulator may affect oogenesis and the development of laid eggs. Hussein *et al.* (2005) proposed that Cry 3Aa protein sequestered by *S. littoralis* binds to ovarian cadherin-like Cry receptors and disturbs the formation of functional egg chambers.

This may explain the reduction in fecundity and fertility after treatment with *B. thuringiensis*. The reduction in fecundity of *S. littoralis* treated with *S. marcescens* might be attributed to the high adult malformation that may prevent males to reach or identify the females for mating process. The higher reduction in the fecundity and fertility in the sequential combined effect treatments

compared to the individual treatments is due to the effect of two stress factors that reflect the synergistic effect of these treatments on *S. littoralis*

Biochemical Effects:

Effects on acid and alkaline phosphatase activities:

Effects on acid phosphatase activity:

Table (4) shows acid phosphatase activity in pupal stage of *S. littoralis*. The obtained results indicated that, in the individual treatments, *S. marcescens* as well as Teflubenzuron exhibited stimulating effect on acid phosphatase activity during pupal stage except the last two times for *S. marcescens* where the activity were reduced to 14.42 % and 74.93%, respectively. On the other hand, individual treatment with *B. thuringiensis* caused significant decrease in acid phosphatase activity during pupal stage except the 2nd and 6th day of pupation, where significant increases were observed in activity comparing to the control. In general, during the pupal stage, the effect of Teflubenzuron was higher than *S. marcescens* which has a higher effect than *B. thuringiensis*.

Table 4: Changes in acid phosphatase activity (µg phenol/min/g body weight) in pupal stage of *Spodoptera littoralis* treated with *Serratia marcescens* at concentration causes 50% malformation, LC₅₀ of *B. thuringiensis*, Teflubenzuron and their combined effects (*Mean ± SE).

Time in day	Acid phosphatase activity (µg phenol/min/g body weight) (*Mean± SE)										
	TREATMENTS										
	Control	<i>Serratia marcescens</i>	***	<i>B. thuringiensis</i>	***	Teflubenzuron	***	<i>B. thuringiensis</i> / <i>Serratia marcescens</i>	***	Teflubenzuron / <i>Serratia marcescens</i>	***
2	11.602 ± 0.305 ^s	20.683 ± 0.337 ^{ijk}	78.27	17.042 ± 0.405 ^{pq}	46.89	27.153 ± 0.983 ^g	134.04	6.673 ± 0.130 ^{uv}	-42.48	40.351 ± 0.784 ^d	247.79
4	21.101 ± 0.615 ^{ij}	45.469 ± 1.303 ^b	115.48	10.060 ± 0.196 st	-52.32	42.315 ± 1.091 ^c	100.54	17.365 ± 0.355 ^{op}	-17.71	55.176 ± 1.330 ^a	161.49
6	10.999 ± 0.155 st	26.867 ± 0.721 ^g	144.27	17.517 ± 0.524 ^{nop}	59.26	23.823 ± 0.456 ^h	116.59	16.318 ± 0.443 ^{pq}	48.36	19.054 ± 0.510 ^{lmn}	73.23
8	20.451 ± 0.342 ^{ijklm}	26.182 ± 0.704 ^g	28.02	10.274 ± 0.152 st	-49.76	35.316 ± 0.643 ^e	72.69	17.429 ± 0.259 ^{op}	-14.78	19.300 ± 0.297 ^{klm}	-5.63
10	16.963 ± 0.403 ^{pq}	19.020 ± 0.288 ^{lmn}	12.13	13.331 ± 0.252 ^r	-21.41	25.758 ± 0.335 ^g	51.85	17.433 ± 0.272 ^{op}	2.77	10.582 ± 0.126 st	-37.62
12	22.092 ± 0.541 ⁱ	18.907 ± 0.421 ^{mno}	-14.42	16.692 ± 0.349 ^{pq}	-24.44	39.996 ± 0.746 ^d	81.04	9.487 ± 0.212 ^t	-57.06	19.602 ± 0.349 ^{ijklm}	-11.27
14	20.501 ± 0.626 ^{ijkl}	5.139 ± 0.076 ^v	-74.93	7.866 ± 0.118 ^u	-61.63	29.166 ± 0.679 ^f	42.27	15.555 ± 0.358 ^q	-24.13	13.858 ± 0.185 ^r	-32.40
LSD	1.5456										

*Means with the same letter(s) are not significantly different.

** The reduction and induction percentage in the enzyme activity with compared to control.

In the sequential combined effect treatments, Teflu /Serr had more stimulating effect by 247.79% and 161.49% at first two times, respectively, than that of both treatments with *S. marcescens* (78.27% and 115.48%, respectively) and Teflubenzuron (134.04% and 100.54%, respectively), whereas, it caused less effect on acid phosphatase activity at the other times of pupation than that of both individual treatments with *S. marcescens* and Teflubenzuron, as compared to control. Regarding to *Bt/Serr*, significant decreases during pupal stage comparing to control were found, whereas, it still less effective as comparing to treatment with *B. thuringiensis*.

Effects on alkaline phosphatase activity:

Table (5) shows changes in the alkaline phosphatase activity during pupal stage of *S. littoralis*. The obtained results revealed that, in the individual treatments, significant decreases from 8th to 12th day of pupation were observed in case of Teflubenzuron treatment which was higher than that of *B. thuringiensis*

whereas, *S. marcescens* had less reducing effect on the alkaline phosphatase activity. also, the individual treatment with Teflubenzuron caused significant increases higher than *S. marcescens* at the first two times of pupation. In the sequential combined effect treatments, Teflu /Serr caused more effect of induction at 2nd day than that of individual treatments either of *S. marcescens* or Teflubenzuron in addition more pronounced reduction at last three times (-76.64%, -91.51% and -86.61%, respectively) than that of *S. marcescens* (-16.14%, -22.17% and -35.68%, respectively) as compared to control. Moreover, *Bt/Serr* behave the same trend of *S. marcescens* except that *Bt/Serr* has more significant elevation effect at 4th and 6th days (109.46 % and 214.93 %, respectively.) as well as more reducing effect at the last three times (-58.16%, -35.51%, and -59.42%, respectively) than that of individual *S. marcescens*, as compared to control.

Table 5: Changes in Alkaline phosphatase activity ($\mu\text{g phenol/min/g body weight}$) in pupal stage of *Spodoptera littoralis* treated with *Serratia marcescens* at concentration causes 50% malformation, LC_{50} of *B. thuringiensis*, Teflubenzuron and their combined effects (*Mean \pm SE).

Time in day	Alkaline phosphatase activity ($\mu\text{g phenol/min/g body weight}$) (*Mean \pm SE)										
	TREATMENTS										
	Control	<i>Serratia marcescens</i>	%	<i>B. thuringiensis</i>	%	Teflubenzuron	%	<i>B. thuringiensis</i> / <i>Serratia marcescens</i>	%	Teflubenzuron / <i>Serratia marcescens</i>	%
2	6.580 \pm 0.199 ^j	16.680 \pm 0.720 ^c	153.50	1.501 \pm 0.034 ^{opq}	-77.19	20.174 \pm 0.322 ^a	206.60	9.874 \pm 0.218 ^{gh}	50.06	16.344 \pm 0.353 ^{ed}	148.39
4	6.224 \pm 0.168 ^j	9.414 \pm 0.148 ^h	51.25	10.171 \pm 0.251 ^g	63.42	15.943 \pm 0.445 ^d	156.15	13.037 \pm 0.221 ^e	109.46	19.488 \pm 0.603 ^b	213.11
6	3.416 \pm 0.067 ^k	6.587 \pm 0.098 ^j	92.83	0.440 \pm 0.028 ^{tw}	-87.12	1.508 \pm 0.034 ^{opq}	-55.85	10.758 \pm 0.203 ^f	214.93	8.515 \pm 0.126 ⁱ	149.27
8	2.536 \pm 0.034 ^{lm}	1.339 \pm 0.034 ^{opqr}	-47.20	1.275 \pm 0.023 ^{opqr}	-49.72	0.996 \pm 0.023 ^{qrst}	-60.73	1.061 \pm 0.023 ^{pqr}	-58.16	2.666 \pm 0.047 ^{lm}	5.13
10	1.605 \pm 0.047 ^{nop}	1.346 \pm 0.017 ^{opqr}	-16.14	0.492 \pm 0.023 ^{stuv}	-69.35	0.233 \pm 0.011 ^v	-85.48	1.035 \pm 0.017 ^{qrs}	-35.51	0.375 \pm 0.028 ^{uv}	-76.64
12	2.743 \pm 0.047 ^l	2.135 \pm 0.022 ^{mn}	-22.17	1.766 \pm 0.030 ^{no}	-35.62	1.326 \pm 0.017 ^{opqr}	-51.66	1.113 \pm 0.017 ^{pqr}	-59.42	0.233 \pm 0.011 ^v	-91.51
14	1.449 \pm 0.069 ^{opqr}	0.932 \pm 0.011 ^{rstu}	-35.68	1.242 \pm 0.022 ^{opqr}	-14.29	0.116 \pm 0.011 ^v	-91.99	2.672 \pm 0.056 ^{lm}	84.40	0.194 \pm 0.022 ^v	-86.61
LSD	0.5574										

*Means with the same letter(s) are not significantly different.

** The reduction and induction percentage in the enzyme activity with compared to control .

The results either of *S. marcescens* or Teflubenzuron were in agreement with those reported by Tolba (2006) when *A. ipsilon* pupae were treated either with *S. marcescens* or Flufenoxuron. On the other hand, the decreased activity in acid phosphatase due to *B. thuringiensis* treatment is similar to Kamel *et al.* (2010) who found decrease in acid phosphatase activity of *S. littoralis* larvae after 48hrs post-treatment with *B. thuringiensis* commercial formulations (Agerin, Dipel 2x and Dipel DF), whereas, it was increased after 120 hrs. for Agerin. The increase in acid and alkaline phosphatase activities during the first period of pupation following treatment with *S. marcescens* may be due to the bacterial growth inside the insect body, this process required more energy and nutrients, so the insect increases acid and alkaline phosphatase activities to compensate the reduction in energy and in tissues development (Tolba 2006). On the other hand, the decrease in alkaline phosphatase activity might be due to the partial lysis of pupal tissues and some of these tissues were retained in the formed adult. Thus, the retention of pupal tissues was a result of the depression in the alkaline phosphatase activity that led to failure in the formation of normal healthy adults (Bassal and Ismail, 1985). Acid and alkaline phosphatases have been shown to be associated with insect development especially in relation to nutrition and egg maturation (Tsumuki and Kanehisa 1984). Acid phosphatase has received considerable attention in developmental studies because of its association with histolysis. This later process is appreciable at the metamorphic moults of holometabolous species to which *S. littoralis* belongs. This may explain the decrease in fecundity and fertility of these compounds and their sequential

combined effects toward *S. littoralis* which obtained in the present study.

Effects on Total Carbohydrate contents:

Table (6) shows changes in total carbohydrate content in pupal stage of *S. littoralis* the obtained results revealed that all treatments cause significant decreases in total carbohydrate content at all times compared to control except the individual treatment with Teflubenzuron in which there were significant increases at 2nd and 10th days, whereas, significant decreases were observed at the other times. The decrements for the individual treatment with *S. marcescens* were gradually pronounced with time reaching drastical decreases at 10th and 12th days of pupation time, which more effective either than *B. thuringiensis* at the time from 4th to 14th days of pupation or Teflubenzuron at 4th, 6th, 8th, 12th and 14th day of pupation time. Moreover, the sequential combined effect treatment, *Bt/Serr*, had higher reducing effect at all times than the individual treatment with *B. thuringiensis* and than the individual *S. marcescens* from 2nd to 10th days of pupation time. On the other hand, the sequential combined effect Teflu /*Serr* had reducing effect at 4th, 6th, 8th, 12th and 14th days of pupation which was higher than those of individual treatment with Teflubenzuron. Thus, it is worth to mention that all treatments contained *S. marcescens* induced more reducing effect on total carbohydrate content of *S. littoralis* pupae than those of Teflubenzuron or *B. thuringiensis* alone. Therefore, it could be stated that *S. marcescens* enhances the reducing effect either of Teflubenzuron or *B. thuringiensis* on the carbohydrates content of *S. littoralis*.

Similar results were reported for total soluble carbohydrate content of pupae of *A. ipsilon* after pupation in sawdust treated with *S. marcescens* or

Flufenoxuron (Tolba, 2006) as well as for total soluble carbohydrate content of 6th instar larvae of *Tribolium castaneum* treated with *Bacillus thuringiensis* (Saleem *et al.*, 1995).

The decrease in total soluble carbohydrate content of untreated pupae

could be attributed to convert it into glucose for supporting all life processes. However, the decrease in total soluble carbohydrate content of treated pupae could be attributed to metamorphic changes in pupa.

Table 6: Changes in total carbohydrates (mg glucose/g body weight) in pupal stage of *Spodoptera littoralis* treated with *Serratia marcescens* at concentration causes 50% malformation, LC₅₀ of *B. thuringiensis*, Teflubenzuron and their combined effects (*Mean ± SE).

Time in day	Total carbohydrates content (µg glucose/g body weight) (*Mean ± SE)										
	TREATMENTS										
	Control	<i>Serratia marcescens</i>	%	<i>B. thuringiensis</i>	%	Teflubenzuron	%	<i>B. thuringiensis</i> / <i>Serratia marcescens</i>	%	Teflubenzuron / <i>Serratia marcescens</i>	%
2	36.634 ± 0.905 ^c	26.522 ± 0.862 ^e	-27.60	22.898 ± 0.843 ^{ghi}	-37.50	45.590 ± 1.271 ^a	24.45	19.635 ± 0.669 ^{hl}	-46.40	24.424 ± 0.703 ^{fg}	-33.33
4	30.169 ± 0.596 ^d	20.514 ± 0.522 ^{jk}	-32.00	31.289 ± 1.423 ^d	3.71	24.615 ± 0.776 ^{efg}	-18.41	16.037 ± 0.562 ^{mn}	-46.84	20.673 ± 0.577 ^{jk}	-31.48
6	36.138 ± 0.700 ^c	21.633 ± 0.623 ^{ij}	-40.14	30.178 ± 0.876 ^d	-16.49	22.300 ± 0.689 ^{hij}	-38.29	15.080 ± 0.480 ^{no}	-58.27	17.907 ± 0.583 ^{lm}	-50.45
8	45.708 ± 0.983 ^a	20.440 ± 0.607 ^{jk}	-55.28	21.445 ± 1.220 ^{ijk}	-53.08	26.183 ± 0.710 ^{ef}	-42.72	11.451 ± 0.381 ^{pqr}	-74.95	18.141 ± 0.504 ^l	-60.31
10	37.191 ± 0.759 ^c	7.092 ± 0.224 ^t	-80.93	24.350 ± 0.927 ^{fg}	-34.53	42.113 ± 1.069 ^b	13.23	6.653 ± 0.278 ^t	-82.11	21.395 ± 0.489 ^{ijk}	-42.47
12	30.886 ± 0.627 ^d	5.693 ± 0.198 ^{tu}	-81.57	24.077 ± 0.783 ^{gh}	-22.05	24.182 ± 0.942 ^{gh}	-21.71	10.423 ± 0.329 ^{qrs}	-66.25	15.364 ± 0.243 ⁿ	-50.26
14	19.612 ± 0.378 ^{hl}	4.098 ± 0.098 ^u	-79.10	11.669 ± 0.434 ^{pq}	-40.50	13.257 ± 0.520 ^{op}	-32.40	9.539 ± 0.374 ^{rs}	-51.36	9.268 ± 0.166 ^s	-52.74
LSD	1.9851										

*Means with the same letter(s) are not significantly different.

** The reduction and induction percentage in the enzyme activity with compared to control .

During this stage, the carbohydrate content supply the body with glucose which provides an energy source for synthesis of pupa and adult tissues, especially the cuticle. Carbohydrates are necessary for the normal functioning of the male and female reproductive systems, as well as for the development of the embryo. In males, sugars form an important constituent of the reproductive glands and most of the carbohydrates of reproductive system were present in the testes. In the females system, carbohydrates are necessary for vitellogenesis and for the formation of the glycosaminoglycans present in the vitelline membrane and chorion. Vitellogenesis involves the accumulation within the oocyte of carbohydrate, lipid and protein yolk to meet the structural

and metabolic needs of the developing embryo (Chippenadale, 1978).

Serratia marcescens caused a high gradual decrease in total carbohydrates in *S. littoralis* pupae because of their requirements to glucose as energy and carbon source for propagation and growth. This may decrease the available carbohydrates in treated insect especially glucose which plays an important role in energy supply, adult maturation (sperm and oocyte development) and builds up a new chitin (Tolba, 2006). This may explain the reduction in fecundity and fertility in the treated insect. Also; this may explain the higher reduction in the fecundity and fertility for sequential combined effect treatments than those of individual treatments either with Teflubenzuron or *B. thuringiensis*.

CONCLUSION

Finally, it could be concluded that *Serratia marcescens* has the potentialities to reduce population density of *S. littoralis* (fertility and fecundity). *Serratia marcescens* also has the ability to increase adult malformation. The obtained results also clearly revealed that the effect of *S. marcescens* on the biological aspect was higher than *B. thuringiensis*, thus, *S. marcescens* can consider a good biopesticide. Also, *S. marcescens* enhances the effect of both *B. thuringiensis* and Teflubenzuron on the biological and biochemical aspects of *S. littoralis*. Therefore, *S. marcescens* could be used either with *B. thuringiensis* or Teflubenzuron through sequential treatments as a good way for the biological control.

REFERANCES

- Abdel-Aal, A. E. and Abdel-khalek, Azza A. (2006). Effect of insect growth regulators on some biological and physiological aspects of *Spodoptera littoralis* (Boisd.). Bull.ent.Egypt,Econ. Ser., 32:101-112.
- Abd El-Hameed, N. F. (1995). Evaluation of the efficacy of some bacterial preparations in controlling the pink bollworm. M.Sc. Thesis, Fac. Agric., Moshtohor (Benha Branch), Zagazig Univ., Egypt, pp: 113-120.
- Abo El-Ghar, G. E. S.; Khalil, M. S. and Eid, T. M. (1994). Effect of plant extracts on development and fecundity of *Agrotis ipsilon* (Lepidoptera : Noctuidae). Bull. ent. Soc. Egypt, Econ. Ser., 21: 171-190.
- Aronson, A.I., Beckman, W. and Dunn, P. (1986) *Bacillus thuringiensis* and related insect pathogens. Microbial. Rev., 50: 1-24.
- Ascher, K.R.S. ; Nemny, N.E. (1984). The effect of CME 134 on *Spodoptera littoralis* eggs and larvae. Phytoparasitica 12: 13-27.
- Bassal, T. T. M. and Ismail, I. E. (1985). Acid and alkaline phosphatases of normal and juvenilized *Spodoptera littoralis* (Noctuidae, Lepidoptera) during metamorphosis. Proc. Zoll. Soc. Egypt, 9: 249-256.
- Brurberg, M. B.; Nes, I. F. and Eijsink, V. G. (2000). Comparative studies of chitinases A and B from *Serratia marcescens*. Microbiology, Jul, 142 (7): 1581-1589.
- Chippenadale, G. M. (1978). The function of carbohydrates in insect life processes. Biochemistry of insects. (Edited by Rockstein, M.) pp. 2-54. Academic press, New York, San Francisco, London.
- Desneux, N. ; Decourtye, A. and Delpuech, J. M. (2007).The sublethal effects of pesticides on beneficial arthropods. Annu. Rev. Entomol. 52: 81-106.
- Dulmage,T. H. and Co-operators (1981). Insecticidal activity of isolates of *Bacillus thuringiensis* and their potential for pest control. In "Microbial Control of Pests and Plant Diseases 1970 - 1980." (H.D. Burges, ed.). pp. 193 -223. Academic Press, London and New York
- Finney, D. J. (1971). Probit Analysis, A statistical treatment of the sigmoid response curve. 7th Ed., Cambridge Univ. Press, Cambridge, England.
- Frank, R.; Braun, H. E.; Ripley, B. D. and Clegy, B. S. (1990). Contamination of rural ponds with pesticides, 1971-1985. Ontario, Canada Bull. Environm. Contamin. Toxicology. 13: 771-817.
- Haseeb, M. and Amano, H. (2002). Effects of contact, oral and persistent toxicity of selected pesticides on *Cotesia plutellae* (Hym., Braconidae), a potential parasitoid of *plutella*

- xylostella (Lep., plutellidae). J. Appl. Entomol. 126: 8-13.
- Hussein, H.M.; Habuštová, O. and Sehnal, F. (2005). Beetle-specific *Bacillus thuringiensis* Cry3Aa toxin reduces larval growth and curbs reproduction in *Spodoptera littoralis* (Boisd.). Pest Manag Sci., 61:1186-1192. DOI: 10.1002/ps.1112.
- Kamel, S.A.; Abd-El Aziz, F.A. and El-Bakry, M.N.(2010). Biochemical effects of three commercial formulations of *Bacillus thuringiensis* (Agerrin, Dipel 2X and Dipel DF) on *Spodoptera littoralis* larvae . Egypt. Acad . J . biolog. Sci.,3 (1): 21-29.
- Laufer, H. and Schin, K. S. (1971). Quantitative studies of hydrolytic enzymes activity in the salivary gland of *Chironomus tentans* (Diptera :Chironomidae) during metamorphosis. Can. Entomol., 103: 454-457.
- Lundgren, L. (1975) Natural plant chemicals acting as oviposition deterrents on cabbage butterflies, *Pieris brassicae* (L.) *P. rapa* (L.) and *P. napi* (L.). Zool. ser., 4: 250 - 258.
- Metcalf, F.; Lu, P. I. and Bowlus, S. (1975). Degradation and environmental fate of 1-(2,6-diflubenzoyl)-3-(4-chlorophenyl) urea. J. Agric. Food Chem., 23: 359-364.
- Saleem, M. A.; Tufail, N. and Shakoori, A. R. (1995). Synergistic effect of synthetic pyrethroids on the toxicity of *Bacillus thuringiensis* as shown by the biochemical changes in the sixth instar larvae of *Tribolium castaneum*. Pakistan J. Zool., 27: (4), 317-323.
- Sikorowski, P. P. (1985). Pecan weevil pathology, pp. 87-101. In W. W. Neel (ed.) Pecan weevil: research perspective. Quail Ridge Press, Jackson, MS.
- Singh, N. B. and Sinha, R. N. (1977): Carbohydrates, lipids and proteins in the developmental stages of *Sitophilus oryzae* and *S. granaries* (Coleoptera: Curculionidae). Ann. Entomol. Soc. Amer., 70: 107-111.
- Sun, Y. P. (1950). Toxicity index-An improved method of comparing the relative toxicity of insecticides. J. Econ. Entomol., 43: 45-53.
- Tolba, H . I. (2006) Biochemical studies on *Serratia marcescens* for controlling the black cutworm, *Agrotis ipsilon* (Huf.) Ph. D. Thesis, Faculty of Agriculture, Cairo University, Egypt. pp. 57-72.
- Trevor, A. J.; Mark, R. M. and Dharmadasa, M. (2004). Use of *Serratia* spp. bacteria for monitoring behaviour of parasitoids. International J. Pest Manage. 50 (3): 173 -176.
- Tsumuki, H. and Kanehisa, K. (1984). Phosphatases in the rice stem borer, *Chilo suppressalis* (Walker) (Lepidoptera: Pyralidae): Some properties and changes of the activities during hibernation. Cryobiology, 21: 177-182.

ARABIC SUMMARY

دراسة مقارنة على التأثيرات البيولوجية والبيوكيميائية لبكتيريا باسليلس ثيورينجينسيز و سيراشيا مرسيسنس و منظم النمو الحشرى تيفلوبينزورون و أيضا تأثيراتهم المشتركة المتعاقبة ضد دودة ورق القطن *سبوتوبتيرا لنيوراليس* (بويدد).

- طارق عفيفي عبد الحميد الشيخ¹ - هبة سمير رافع² - عبد المنعم محمد الاعسر³ - صفوت حسن علي⁴
- 1- معهد بحوث وقاية النباتات - مركز البحوث الزراعية الدقي الجيزة
 - 2- معهد بحوث وقاية النباتات - مركز البحوث الزراعية الدقي الجيزة
 - 3- قسم الكيمياء الحيوية - كلية الزراعة - جامعة عين شمس
 - 4- قسم الكيمياء الحيوية - كلية الزراعة - جامعة عين شمس

اشتمل البحث على دراسة بعض التأثيرات البيولوجية و البيوكيميائية الناتجة عن استخدام التركيز المتسبب في تشوه 50% من الفراشات لبكتيريا سيراشيا مرسيسنس المعامل بها في التربه و التركيز القاتل للنصف لكلا من المستحضر التجارى بروتيكنو (بكتيريا باسليلس ثيورينجينسيز) و منظم النمو الحشرى تيفلوبينزورون ضد العمر الثانى لدودة ورق القطن و ايضا تأثيراتهم المشتركة من خلال معاملة العمر الثانى بالتركيز القاتل للنصف لكلا من الباسلليس ثيورينجينسيز و التيفلوبينزورون ثم تعريض اليرقات عند التعذير لتربه معامله ببكتيريا سيراشيا مرسيسنس عند التركيز المتسبب فى تشوه 50% من الفراشات. أظهرت كل المعاملات نقص معنوى فى نسب وضع و فقس البيض و الذى كان أكثر وضوحا فى معاملات التأثيرات المشتركة . تسببت المعاملات فى تأثيرات معنويه على نشاط انزيمات الفوسفاتيز القلوي و الحامضي و المحتوى الكلى للكربوهيدرات للعدارى فى الفترات المختلفة للطور العذري . حيث تسببت المعامله بكلا من التيفلوبينزورون و السيراشيا مرسيسنس الى زيادة نشاط إنزيم الفوسفاتيز الحامضي على مدار فترات الطور العذرى ما عدا اخر فترتين فى المعامله بالسيراشيا مرسيسنس حيث حدث انخفاض معنوى فى نشاط إنزيم الفوسفاتيز الحامضي بينما ادت السيراشيا مرسيسنس المعامله فى التربه بعد معاملة العمر الثانى بالتيفلوبينزورون الى زياده معنويه فى نشاط إنزيم الفوسفاتيز الحامضي على مدار الثلاث فترات الاولى ثم انخفاض النشاط بعد ذلك حتى نهاية الطور العذرى. بينما تسببت المعاملات الاخرى ايضا فى انخفاض معنوى فى نشاط إنزيم الفوسفاتيز الحامضي فى معظم فترات الطور العذرى . أدت السيراشيا مرسيسنس سواء المعاملة فرديا او بعد كلا من معاملات الباسلليس ثيورينجينسيز أو التيفلوبينزورون الى زياده معنويه فى نشاط إنزيم الفوسفاتيز القلوى على مدار الثلاث فترات الاولى ثم انخفاض النشاط بعد ذلك حتى نهاية الطور العذرى . بينما أدت المعامله بالباسلليس ثيورينجينسيز الى انخفاض معنوى على مدار الطور العذرى . ايضا تسببت كل المعاملات فى انخفاض معنوى فى المحتوى الكلى للكربوهيدرات فى معظم فترات الطور العذرى. و النتائج التى تم الحصول عليها تشير الى أن بكتيريا سيراشيا مرسيسنس يمكن اعتبارها مبيد حيوى وتكون أيضا أكثر تأثيرا فى المعاملات المشتركة بالتتابع مع الباسلليس ثيورينجينسيز أو التيفلوبينزورون .